



| |
|---------|
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |
| Level 1 |



| |
|------|
| 4843 |
| 4844 |
| 4845 |
| 4846 |
| 4847 |
| 4848 |
| 4849 |
| 4850 |
| 4851 |
| 4852 |
| 4853 |
| 4854 |
| 4855 |
| 4856 |
| 4857 |
| 4858 |
| 4859 |
| 4860 |
| 4861 |
| 4862 |
| 4863 |
| 4864 |

| |
|--|
| N. Carmona-Gutierrez Minois. Spermidine promotes stress resistance in <i>Drosophila melanogaster</i> through autophagy-dependent and -independent pathways. <i>Cell Death Dis.</i> 2012. 3:e401 |
| A. Plyusnina Moskalev. The role of D-GADD45 in oxidative, thermal and genotoxic stress resistance. <i>Cell Cycle.</i> 2012. 11:4222-41 |
| A. G. Harper Jimenez. Linkages between the life-history evolution of tropical and temperate birds and the resistance of cultured skin fibroblasts to oxidative and non-oxidative chemical injury. <i>J Exp Biol.</i> 2013. 216:1373-80 |
| Y. Peng Zuo. Sesamin extends the mean lifespan of fruit flies. <i>Biogerontology.</i> 2013. 14:107-19 |
| A. C. Loh Costa. <i>Drosophila</i> Trap1 protects against mitochondrial dysfunction in a PINK1/parkin model of Parkinson's disease. <i>Cell Death Dis.</i> 2013. 4:e467 |
| E. B. Wisman Menshchikova. ARE-inducing phenol antioxidant TC-13 improves survival of <i>Drosophila melanogaster</i> in oxidative stress. <i>Bull Exp Biol Med.</i> 2012. 154:260-4 |
| C. B. Copes Edwards. Malate and fumarate extend lifespan in <i>Caenorhabditis elegans</i> . <i>PLoS One.</i> 2013. 8:e58345 |
| C. Li Ren. GLB-13 is associated with oxidative stress resistance in <i>Caenorhabditis elegans</i> . <i>IUBMB Life.</i> 2013. 65:423-34 |
| M. Ranji Rauthan. The mitochondrial unfolded protein response activator ATFS-1 protects cells from inhibition of the mevalonate pathway. <i>Proc Natl Acad Sci U S A.</i> 2013. 110:5981-6 |
| R. Hosamani. Acute exposure of <i>Drosophila melanogaster</i> to paraquat causes oxidative stress and mitochondrial dysfunction. <i>Arch Insect Biochem Physiol.</i> 2013. 83:25-40 |
| S. Mukai Bahadorani. Expression of zinc-deficient human superoxide dismutase in <i>Drosophila</i> neurons produces a locomotor defect linked to mitochondrial dysfunction. <i>Neurobiol Aging.</i> 2013. 34:2322-30 |
| R. J. Reiter. Aging and oxygen toxicity: Relation to changes in melatonin. <i>Age (Omaha).</i> 1997. 20:201-13 |
| J. Mu Yang. Changes in the expression of four heat shock proteins during the aging process in <i>Brachionus calyciflorus</i> (rotifera). <i>Cell Stress Chaperones.</i> 2014. 19:33-52 |
| A. Greenough Southon. Presenilin promotes dietary copper uptake. <i>PLoS One.</i> 2013. 8:e62811 |
| S. H. Qamar Mehdi. Paraquat-induced ultrastructural changes and DNA damage in the nervous system is mediated via oxidative-stress-induced cytotoxicity in <i>Drosophila melanogaster</i> . <i>Toxicol Sci.</i> 2013. 134:355-65 |
| J. Feng Zhi de. Realgar bioleaching solution suppress ras excessive activation by increasing ROS in <i>Caenorhabditis elegans</i> . <i>Arch Pharm Res.</i> 2014. 37:390-8 |
| E. M. Lee Vayndorf. Whole apple extracts increase lifespan, healthspan and resistance to stress in <i>Caenorhabditis elegans</i> . <i>J Funct Foods.</i> 2013. 5:1236-1243 |
| J. Gontier Xu. Longevity effect of IGF-1R(+/-) mutation depends on genetic background-specific receptor activation. <i>Aging Cell.</i> 2014. 13:19-28 |
| T. Nemoto-Sasaki Takeuchi. Galectin LEC-1 plays a defensive role against damage due to oxidative stress in <i>Caenorhabditis elegans</i> . <i>J Biochem.</i> 2013. 154:455-64 |
| V. Turm Pandey. A new in vivo model of pantothenate kinase-associated neurodegeneration reveals a surprising role for transcriptional regulation in pathogenesis. <i>Front Cell Neurosci.</i> 2013. 7:146 |
| J. P. Kuiper Melis. Slow accumulation of mutations in Xpc-/- mice upon induction of oxidative stress. <i>DNA Repair (Amst).</i> 2013. 12:1081-6 |
| B. Li Qabazard. Hydrogen sulfide is an endogenous regulator of aging in <i>Caenorhabditis elegans</i> . <i>Antioxid Redox Signal.</i> 2014. 20:2621-30 |

| |
|--|
| The naturally occurring polyamine spermidine (Spd) has recently been shown to promote longevity across species in an a |
| There is a relationship between various cellular stress factors and aging. In earlier studies, we demonstrated that overex |
| A fundamental challenge facing physiological ecologists is to understand how variation in life history at the whole-organi |
| The present study investigated the anti-ageing activity of sesamin and its effect on gene expression of superoxide dismut |
| Mitochondrial dysfunction caused by protein aggregation has been shown to have an important role in neurological dise |
| The effects of hydrophilic synthetic antioxidant TC-13 sodium (3'-(3'-tert-butyl-4'-hydroxyphenyl)propylthiosulfonate on |
| Malate, the tricarboxylic acid (TCA) cycle metabolite, increased lifespan and thermotolerance in the nematode <i>C. elegans</i> |
| Globins constitute a superfamily of heme-binding proteins that is widely present in many species. There are 33 putative g |
| Statins are cholesterol-lowering drugs that inhibit 3-hydroxy-3-methyl-glutaryl-CoA (HMG-CoA) reductase, the rate-limiti |
| Paraquat (PQ; 1, 1'-dimethyl-4-4'-bipyridinium), an herbicide and model neurotoxicant, is identified to be one of the prim |
| More than 130 different mutations in the Cu/Zn superoxide dismutase (SOD1) gene have been associated with amyotrop |
| Melatonin (N-acetyl-5-methoxytryptamine) is a chemical mediator produced in the pineal gland and other sites in the bo |
| Heat shock proteins (HSPs) are molecular chaperones and have an important role in the refolding and degradation of mis |
| Dietary copper is essential for multicellular organisms. Copper is redox active and required as a cofactor for enzymes suc |
| Paraquat (PQ), a quaternary nitrogen herbicide, is commonly used as a pesticide despite of its high toxicity. Our study ev |
| Although realgar bioleaching solution (RBS) has been proved to be a potential candidate for cancer therapy, the mechani |
| Regular consumption of fruits and vegetables is associated with reduced risk of age-related functional decline and chroni |
| Growth hormone (GH) and insulin-like growth factor (IGF) signaling regulates lifespan in mice. The modulating effects of |
| LEC-1 is a major galectin in <i>Caenorhabditis elegans</i> and contains two carbohydrate recognition domains (CRDs), N-CRD at |
| Pantothenate Kinase-Associated Neurodegeneration (PKAN) is a neurodegenerative disorder with a poorly understood m |
| XPC is one of the key DNA damage recognition proteins in the global genome repair route of the nucleotide excision repa |
| AIMS: To investigate the role of endogenous hydrogen sulfide (H ₂ S) in the control of aging and healthspan of <i>Caenorhab</i> |



| |
|------|
| 4865 |
| 4866 |
| 4867 |
| 4868 |
| 4869 |
| 4870 |
| 4871 |
| 4872 |
| 4873 |
| 4874 |
| 4875 |
| 4876 |
| 4877 |
| 4879 |
| 4880 |
| 4881 |
| 4882 |
| 4883 |
| 4884 |
| 4885 |
| 4886 |

| |
|---|
| S. R. Haddadi Jahromi. Neuroprotective effect of <i>Decalepis hamiltonii</i> in paraquat-induced neurotoxicity in <i>Drosophila melanogaster</i> : biochemical and behavioral evidences. <i>Neurochem Res.</i> 2013. 38:2616-24 |
| N. Austad Elbourkadi. Fibroblasts from long-lived species of mammals and birds show delayed, but prolonged, phosphorylation of ERK. <i>Aging Cell.</i> 2014. 13:283-91 |
| W. S. Chen Lin. The anti-aging effects of <i>Ludwigia octovalvis</i> on <i>Drosophila melanogaster</i> and SAMP8 mice. <i>Age (Dordr).</i> 2014. 36:689-703 |
| M. Ganfornina Ruiz. Ligand binding-dependent functions of the lipocalin NLaz: an in vivo study in <i>Drosophila</i> . <i>Faseb j.</i> 2014. 28:1555-67 |
| A. Morales Lucas. Differential effects of specific carotenoids on oxidative damage and immune response of gull chicks. <i>J Exp Biol.</i> 2014. 217:1253-62 |
| H. F. Jimenez-Del-Rio Ortega-Arellano. Dmp53, basket and drICE gene knockdown and polyphenol gallic acid increase life span and locomotor activity in a <i>Drosophila</i> Parkinson's disease model. <i>Genet Mol Biol.</i> 2013. 36:608-15 |
| D. Cha Ahn. The Longevity Properties of 1,2,3,4,6-Penta-O-Galloyl-beta-D-Glucose from <i>Curcuma longa</i> in <i>Caenorhabditis elegans</i> . <i>Biomol Ther (Seoul).</i> 2013. 21:442-6 |
| Y. Zhang-Akiyama Sanada. An increase of oxidised nucleotides activates DNA damage checkpoint pathway that regulates post-embryonic development in <i>Caenorhabditis elegans</i> . <i>Mutagenesis.</i> 2014. 29:107-14 |
| W. Fang Zhao. Nar1 deficiency results in shortened lifespan and sensitivity to paraquat that is rescued by increased expression of mitochondrial superoxide dismutase. <i>Mech Ageing Dev.</i> 2014. 138:53-8 |
| Y. Komura Yaguchi. Influence of oral supplementation with sesamin on longevity of <i>Caenorhabditis elegans</i> and the host defense. <i>Eur J Nutr.</i> 2014. 53:1659-68 |
| W. G. Xavier Salgueiro. Direct synthesis of 4-organylsulphenyl-7-chloro quinolines and their toxicological and pharmacological activities in <i>Caenorhabditis elegans</i> . <i>Eur J Med Chem.</i> 2014. 75:448-59 |
| R. Kumimoto Islam. ALS-linked SOD1 in glial cells enhances ss-N-Methylamino L-Alanine (BMAA)-induced toxicity in <i>Drosophila</i> . <i>F1000Res.</i> 2012. 1:47 |
| Y. Asai Sanada. Oxidation resistance 1 is essential for protection against oxidative stress and participates in the regulation of aging in <i>Caenorhabditis elegans</i> . <i>Free Radic Res.</i> 2014. 48:919-28 |
| U. Yu Na. The LYR factors SDHAF1 and SDHAF3 mediate maturation of the iron-sulfur subunit of succinate dehydrogenase. <i>Cell Metab.</i> 2014. 20:253-66 |
| T. B. Ida Nguyen. Role of SCOX in determination of <i>Drosophila melanogaster</i> lifespan. <i>Am J Cancer Res.</i> 2014. 4:325-36 |
| A. M. Lehr Pickering. Fibroblasts From Longer-Lived Species of Primates, Rodents, Bats, Carnivores, and Birds Resist Protein Damage. <i>J Gerontol A Biol Sci Med Sci.</i> 2015. 70:791-9 |
| M. Issa Cassar. A dopamine receptor contributes to paraquat-induced neurotoxicity in <i>Drosophila</i> . <i>Hum Mol Genet.</i> 2015. 24:197-212 |
| W. S. Ludwig Chick. Screening for stress-resistance mutations in the mouse. <i>Front Genet.</i> 2014. 5:310 |
| A. B. Ryu Hwang. Feedback regulation via AMPK and HIF-1 mediates ROS-dependent longevity in <i>Caenorhabditis elegans</i> . <i>Proc Natl Acad Sci U S A.</i> 2014. 111:E4458-67 |
| A. K. Ratnasekhar Shukla. Metabolomic Analysis Provides Insights on Paraquat-Induced Parkinson-Like Symptoms in <i>Drosophila melanogaster</i> . <i>Mol Neurobiol.</i> 2016. 53:254-69 |
| Y. Lin Sun. Sterols from Mytilidae show anti-aging and neuroprotective effects via anti-oxidative activity. <i>Int J Mol Sci.</i> 2014. 15:21660-73 |

| |
|---|
| In this paper, we have demonstrated for the first time, the antioxidant and neuroprotective effects of <i>Decalepis hamiltonii</i> |
| Fibroblasts from long-lived mutant mice show diminished phosphorylation of the stress-activated protein kinases ERK1/2 |
| We investigated the anti-aging effects of <i>Ludwigia octovalvis</i> (Jacq.) P. H. Raven (Onagraceae), an extract of which is widely |
| Lipocalins are small extracellular proteins mostly described as lipid carriers. The <i>Drosophila</i> lipocalin NLaz (neural Lazarin) |
| Micronutrients are essential for normal metabolic processes during early development. Specifically, it has been suggested |
| Understanding the mechanism(s) by which dopaminergic (DAergic) neurons are eroded in Parkinson's disease (PD) is critical |
| Here in this study, we isolated 1,2,3,4,6-penta-O-galloyl-beta-D-glucose (PGG) from <i>Curcuma longa</i> L. and elucidated the |
| 8-Oxo-dGTP, an oxidised form of dGTP generated in the nucleotide pool, can be incorporated opposite adenine or cytosine |
| <i>Saccharomyces cerevisiae</i> Nar1p is an essential Fe/S protein that exhibits striking similarity to bacterial iron-only hydrogenase |
| PURPOSE: Nutritional control has been proposed as a potential therapy for slowing the senescence of immune function and |
| We describe herein our results on the synthesis and biological properties in <i>Caenorhabditis elegans</i> of a range of 4-organ |
| Environmental factors have been implicated in the etiology of a number of neurodegenerative diseases, including amyotrophic |
| Human oxidation resistance 1 (OXR1) functions in protection against oxidative damage and its homologs are highly conserved |
| Disorders arising from impaired assembly of succinate dehydrogenase (SDH) result in a myriad of pathologies, consistent |
| In man, COX (cytochrome c oxidase) deficiency is reported to be related to mutation of the SCO2 (synthesis of cytochrome |
| Species differ greatly in their rates of aging. Among mammalian species life span ranges from 2 to over 60 years. Here, we |
| Long-term exposure to environmental oxidative stressors, like the herbicide paraquat (PQ), has been linked to the development |
| Longevity is correlated with stress resistance in many animal models. However, previous efforts through the boosting of |
| Mild inhibition of mitochondrial respiration extends the lifespan of many species. In <i>Caenorhabditis elegans</i> , reactive oxygen |
| Paraquat (PQ) exposure causes degeneration of the dopaminergic neurons in an exposed organism while altered metabolic |
| For screening anti-aging samples from marine natural products, K6001 yeast strain was employed as a bioassay system. The |

| |
|--------------|
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |
| Not Relevant |